

CHARLES UNIVERSITY IN PRAGUE
FACULTY OF NATURAL SCIENCES



BACHELOR THESIS

Cyclophosphamide Effects on Hematopoiesis

Účinky cyklofosfamidu na hematopoesu

Nicol Renešová

Supervisor: RNDr. Luděk Šefc, CSc.

Prague 2011

Author's declaration:

I declare that I have written this thesis independently with the help of literature and sources listed. Neither this thesis nor any substantial part of it was submitted with the aim to obtain another or the same academic degree.

17th August 2011, Prague

Acknowledgments:

I wish to thank my supervisor RNDr. Luděk Šefc, CSc. for valuable advice concerning the thesis and its structure, and my family and friends who supported me throughout the process of its creation.

Abstract

Cyclophosphamide is an alkylating agent developed in the 1960s for the use in treatment of cancerous diseases. Since its introduction, it has manifested various spectra of effects. Of uttermost importance are the impacts cyclophosphamide has on the hematopoietic system housed in the bone marrow of femoral and other bones. Hematopoiesis is a complex process which has been extensively studied for decades. The more it is known about the niches the hematopoietic stem cells occupy as well as of their life cycle, the more it is possible to design functional therapy for its malignancies and improve the survival rates. The effects of cyclophosphamide administration on hematopoietic system are divided into three major categories: myeloablation and myelosuppression, immunosuppression, and mobilisation of hematopoietic stem cells into the peripheral blood. The immunosuppression is achieved by systematic depletion of progenitors differentiating into leucocytes. Induced neutropenia and lymphopenia allow for management of autoimmune diseases and preservation of transplants. Mobilisation, a process opposite to stem cell homing, seems to be dependent on disruption of cellular adhesion (CXCR4/SDF-1, VCAM-1/VLA-4) facilitated by proteases released into the environment after cyclophosphamide exposure.

Key words: cyclophosphamide, hematopoietic stem cells, cytochrome P450, cytochrome P450 reductase, immunosuppression, mobilisation, cellular adhesion.

Abstrakt

Cyklofosfamid je alkylační činidlo vyrobené v šedesátých letech minulého století za účelem použití v protirakovinné terapii. Od svého uvedení manifestoval široké spektrum účinků. Velkou pozornost si zaslouží zejména účinky, které cyklofosfamid má na krvetvorbu, jež je lokalizována v dřeni stehenních a dalších kostí. Krvetvorba je složitý proces, jehož výzkum se odehrává už řadu desetiletí. Čím více je toho známo o takzvaných nikách okupovaných hematopoetickými kmenovými buňkami, jakožto i o jejich životním cyklu, tím snadněji se dá vyvíjet účinná terapie proti malignitám, což ve výsledku vede i k navýšení počtu přeživších pacientů. Dopad cyklofosfamidu na krvetvorbu se dá shrnout třemi hlavními pojmy: myeloablace a myelosuprese, imunosuprese, mobilizace hematopoetických kmenových buněk do periferní krve. Imunosuprese je dosaženo systematickou deplecí progenitorových buněk diferencujících se do bílých krvinek. Navozená neutropenie a lymfopenie umožňují zmírnění (i když většinou nikoliv vyléčení) autoimunitních chorob a zabránění odmítnutí štěpu po allotransplantaci. Mobilizace, proces opačný k takzvanému „homingu“ krvetvorných kmenových buněk, se zdá být závislou především na eliminaci buněčné adheze (CXCR4/SDF-1; VCAM-1/VLA-4) prostřednictvím proteáz vyloučených do prostředí niky po vystavení cyklofosfamidu.

klíčová slova: cyklofosfamid, hematopoetické kmenové buňky, cytochrom P450, cytochrom P450 reduktáza, imunosuprese, mobilizace, buněčná adheze.

LIST OF CONTENT

1. Introduction	2
2. Cyclophosphamide Metabolism	3
2.1. Key Enzymes	3
2.1.1...Cytochrome P450	3
2.1.2. CYP Reductase	5
2.2. Activation Pathway	7
2.2.1. 4-hydroxycyclophosphamide	7
2.2.2. Spontaneous Reaction of Aldophosphamide	8
2.2.3. Acrolein	8
2.2.4...Phosphoramidate mustard	9
3. Hematopoiesis	12
3.1. Cyclophosphamide Effects on Hematopoiesis	15
3.1.1. Myeloablation, Myelosuppression and Immunosuppression	15
3.1.2. Mobilisation of Hematopoietic Stem Cells	18
4. Conclusion	21
5. References	22

1. Introduction

N,N-bis(2-chloroethyl)-1,3,2-oxazaphosphinan-2-amine 2-oxide (cyclophosphamide) has been used in humane as well as veterinary medicine for great many years (Henry et al, 1994). It is a potent drug belonging into a family of alkylating agents called oxazaphosphorines that have numerous effects. They are: mutagenic, teratogenic, carcinogenic, immunosuppressive and antineoplastic to name just a few (Kenney et al, 2001). Cyclophosphamide is currently in use worldwide mainly in the treatment of several kinds of cancerous diseases, usually in combination with several other chemotherapeutic agents. It is quite versatile as far as its use in the clinical setting is concerned.

Among its most important effects are those cyclophosphamide has on hematopoiesis. Not only is it able to cause myeloablation and immunosuppression, it is also capable of mobilising hematopoietic stem cells into the peripheral blood and thus enables their use for autotransplantation in patients with severe hematologic malignancies and immune malignancies.

This thesis aims to explore and put into perspective the knowledge that has been accumulated and refined since the first preparation of cyclophosphamide some sixty years ago.

2. Cyclophosphamide Metabolism

Cyclophosphamide in itself is a harmless substance (a prodrug) that needs to be bioactivated to acquire its cytotoxic properties (see Figures 4 and 5).

2.1. Key Enzymes

2.1.1. Cytochrome P450

Cytochrome P450 (CYP) is a name given to a group of enzymes belonging to the class of oxidoreductases. The P450 part of the name is derived from the fact that during a spectroscopic analysis they demonstrate the highest absorbance when the analytic beam of light has the wavelength of 450 nanometres.

These enzymes can be found throughout eukaryotic organisms and in some bacteria (generally those, which use hydrocarbons as a source of carbon as well as energy). In mammals they are included in a wide spectrum of metabolic pathways – steroidogenesis, metabolism of vitamins A and D, eicosanoid metabolism, transformation of xenobiotics (de Montellano, 2005). Some of them possess several functions and respond to the presence of a wider range of substrates. Only a handful of human cytochrome P450s are capable of transformation of xenobiotics, but all of those are capable of processing an endogenous substrate, albeit usually with lower specificity and/or metabolic rate.

CYPs are cysteinato-heme proteins (see Figure 1) which in mammals vary in expression according to the species, localisation, environment – some CYPs are expressed constitutively, others are inducible – age and even gender of the individual. The majority of CYPs are in occurrence in the liver. They are membrane-bound proteins which occur predominantly in the endoplasmic reticulum, even though a small enclave can be encountered in the mitochondria.




Figure 1. Prosthetic of cysteinato-heme enzymes: an iron(III) protoporphyrin-IX linked with a proximal cysteine ligand. (reproduced from Meunier et al, 2004)

Cytochrome P450s have been identified as a crucial component of the cyclophosphamide metabolic pathway, as it is these enzymes that enable cyclophosphamide's bioactivation (Pass et al, 2005; Ren et al, 1997). Cyclophosphamide in the liver is activated by a group of CYP isoenzymes by means of hydroxylation on its C-4 atom. This reaction yields 4-hydroxycyclophosphamide and its tautomeric form aldophosphamide, which are at equilibrium.

CYPs are often referred to as monooxygenases because they use molecular oxygen in the oxidation reactions they catalyse, and incorporate only one of its atoms into the substrate that is being catalysed. The other oxygen atom is reduced to water with two electrons provided by NAD(P)H via a reductase (Equation 1).



Equation 1. Monooxygenation catalysed by a cytochrome P450 enzyme.

The enzymes thought to possess the greatest potential with regards to cyclophosphamide metabolism have been identified as: CYP2B6, CYP2A6, CYP2C9 and CYP3A4 (Roy et al, 1999). While the majority of the applied dose of cyclophosphamide follows the route of 4-hydroxylation and activation, a smaller portion can follow another metabolic pathway – N-dechloroethylation and subsequent formation of inactive and neurotoxic chloroacetaldehyde. CYP3A4 proved to be the most prolific in this alternative metabolism (Huang et al, 2000).

Unfortunately, there is a substantial genetic polymorphism in CYPs, leading to several abnormalities in the metabolism of cyclophosphamide – such as lower substrate specificity, a lower metabolic rate – in some individuals, which may result in the patient being less responsive to treatment by the drug, and other adverse situations (Griskevicius et al, 2003; Xie et al, 2003).

Moreover, cyclophosphamide is often used in combination chemotherapy during which several chemicals are co-administered, therefore possible drug-drug interactions have to be carefully evaluated. For example triethylenethiophosphoramide (thioTEPA) – a common antineoplastic agent – is responsible for specific uncompetitive inhibition of CYP 2B6, and thus interferes with the conversion of cyclophosphamide to 4-hydroxycyclophosphamide (Harleton et al, 2004; Rae et al, 2002). Even cyclophosphamide itself regulates its own metabolism by induction of CYPs with subsequent increase in 4-hydroxylation rate (Chang et al, 1997); on the other hand acrolein that is produced further along the activation pathway is thought to be responsible for inhibiting cyclophosphamide metabolism by interacting with both CYPs and the CYP reductase (LeBlanc & Waxman, 1990).

2.1.2. CYP Reductase

Cytochrome P450 reductase (also known as cytochrome c reductase) is a protein localised on the cytosolic face of the endoplasmic reticulum, and is most abundant in hepatocytes (Phillips & Langdon, 1962). It is safely anchored into the membrane of the endoplasmic reticulum by a sequence of about sixty amino acids. The cytosolic part is comprised of two domains. The first one has two binding sites for NADPH and uses a FAD molecule as a cofactor, while the other one employs a

FMN (flavin mononucleotide) molecule as its cofactor and is responsible for the contact with a CYP molecule. The tertiary structure of the enzyme is highly conserved, indicating its importance.

The reductase shuttles electrons to the CYP in several steps. Upon binding NADPH the FAD-domain is reduced by two hydride ions simultaneously. In the next step, the electrons are channelled to the FMN-domain in sequence, and finally to the CYP (Vermilion et al, 1981). Two conformational states of the reductase have been described – the “open” state is compatible with FMN-heme electron transfer, while the “closed” state allows FAD-FMN electron transfer (Fig. 2) (Ellis et al, 2009).




Figure 2. The open (left) and closed (right) conformations of the CYP reductase. The FMN domain is shown in blue, the linker domain in purple and the FAD domain in green colour. The cofactors are shown: FMN – yellow, FAD – red, NADPH – blue (reproduced from Ellis et al, 2009).

Despite having two binding sites for NADPH, only one of them is catalytic. Simultaneous binding of two molecules of NADPH actually attenuates the reaction,

most probably by interfering with the release of NADP⁺ from the catalytic site (de Montellano, 2005).

The diflavin reductase is vital for monooxygenation reactions of CYPs because it provides the electrons necessary for reduction of the CYP in question, and subsequently the reduction of an oxygen atom to water. An experimental knockout mouse strain deficient in expression of hepatic CYP reductase (Hepatic Cytochrome P450 Null – HNR) – to the extent of expressing less than 1 % of the amount of the enzyme produced by wild-type mice – in which has the conversion of cyclophosphamide to 4-hydroxyphosphamide been significantly halted, and its half-life in plasma dramatically increased, have been used to study the importance of the CYP reductase for CYP-facilitated drug metabolism (Pass et al, 2005). Despite having virtually no hepatic CYP reductase activity, the bioactivation of cyclophosphamide still takes place owing to extra-hepatic activation, but the amount of 4-hydroxycyclophosphamide produced is correspondingly lowered. The extra-hepatic bioactivation of cyclophosphamide has been demonstrated on another mouse strain – the CL-LCN (CYP reductase-low, liver- CYP reductase-null) mice (Gu et al, 2007).

2.2. Activation Pathway

2.2.1. 4-hydroxycyclophosphamide

A tautomer of aldophosphamide, 4-hydroxyphosphamide has been shown to possess little cytotoxic potential and is therefore not considered to be a therapeutic metabolite of cyclophosphamide. Nevertheless, it does have an important function as this is the form in which the drug enters target cells, and tautomerises to aldophosphamide, which then undergoes further metabolic processing leading to the final therapeutic product. It is presumed that 4-hydroxycyclophosphamide travels to its final destination bound to the red blood cells. This notion is largely supported by the findings of higher concentrations of activated oxazaphosphorines in the erythrocytes than in the plasma (Highley et al, 1997).

2.2.2. Spontaneous Reaction of Aldophosphamide

Aldophosphamide that is transported to target tissues by the blood circulatory system undergoes a non-enzymatic elimination to form phosphoramidate mustard and acrolein, former of which is an alkylating agent that forms intra- as well as interstrand cross-links in the DNA and protein-DNA cross-links. Such an intervention to the integrity of the DNA constitutes a significant damage to the cell and may lead to its death by either apoptosis or necrosis – that is the basic principle of cyclophosphamide and related substances' (ifosfamide, trofosfamide) therapeutic effect. On the other hand, acrolein is a highly nephrotoxic and neurotoxic by-product of the reaction. Cytotoxic properties of the compound cause a disease of the urinary bladder called haemorrhagic cystitis (Lawrence et al, 1975).

Even though the reaction does not require a catalyst, it has been established that human serum albumin speeds up the process (Kwon et al, 1987). However, the phosphoramidate mustard generated in the blood does not pose a significant threat to tissues as it is mostly ionised (at physiological pH) which is the underlying reason for its decreased ability to pass through the plasma membrane of cells.

In addition to the spontaneous reaction aldophosphamide can also serve as a substrate for an aldehyde dehydrogenase which catalyses its oxidation to the inactive carboxyphosphamide by molecular oxygen. At the same time a superoxide anion belonging to the group of reactive oxygen species (ROS) is generated. It has been suggested that upregulation of the enzyme contributes to cyclophosphamide resistance (Manthey et al, 1990).

2.2.3. Acrolein

In spite of contributing to the cytotoxic properties of cyclophosphamide, the generation of acrolein quantitatively equal to phosphoramidate mustard is mostly undesirable due to its role in the development of haemorrhagic cystitis and neurotoxicity. Acrolein produced by β -elimination from aldophosphamide causes single-strand breaks in the DNA and readily binds to cellular proteins (Crook et al, 1986). From the fact that it has been known to deplete the cellular glutathione has been derived the practice of co-administration of protective drugs which include

sulphydryl-containing compounds such as mesna (Figure 3), which allow for its safe elimination from the body.

Acrolein is at least on order of magnitude more effective in depletion of glutathione than phosphoramidate mustard, therefore agents responsible for protection against acrolein toxicity can be used without halting the curative effect of cyclophosphamide (Gurtoo et al, 1981).




Figure 3. The structure of mesna.

2.2.4. Phosphoramidate Mustard

The phosphoramidate mustard's potential to cause damage to cells by alkylation of their nucleic acids lies within its electrophilic aziridinium group which naturally seeks a nucleophilic partner to react with. The N-7 of guanines has proven to be a favoured site for alkylation, especially if they are flanked by more guanines. As the mustard possesses two functional groups, it interconnects two reaction sites, thus forming the inter- and intrastrand (most often between directly adjacent bases) crosslinks as well as nucleic acid-protein crosslinks (Kohn et al, 1987).

Cyclophosphamide metabolism has been extensively studied in mice. There is a major difference between human and murine metabolisms that has to be accounted for when applying murine data into clinical practice. While there are no active metabolites detectable in murine blood as soon as one hour after dose administration (personal communication with RNDr. Luděk Šefc, CSc.), the build-up of active metabolites and clearance in humans is slower and happens in a space of several hours. Importantly, there does not seem to be a difference between oral and intravenous routes of administration, which means patients can be offered the more comfortable oral treatment without compromising the therapeutic outcome (Struck et al, 1987).

Cyclophosphamide and its metabolites are mostly cleared from the body via urine.




Figure 4. Metabolism of cyclophosphamide. Vertically: activation of cyclophosphamide; horizontally: inactivation pathways (reproduced from de Jonge et al, 2005).

3. Hematopoiesis

Hematopoiesis is a process during which hematopoietic stem cells (HSCs) maintain their numbers, and produce a cohort of terminally differentiated mature blood cells. During the foetal development hematopoietic activity can be observed in the liver as well as the spleen, but the bone marrow is the only site that supports hematopoiesis after the birth and throughout adulthood. Indeed, extramedullary (occurring outside the bone marrow compartment) hematopoiesis is widely recognised as a sign of a disease and/or a direct consequence of chemotherapeutically or by irradiation impaired hematopoietic capacity of the bone marrow. In these cases, extramedullary hematopoiesis serves as a mechanism for compensation of the lost medullary hematopoiesis (Singer et al, 2004). Splenectomised mice recover more slowly than controls after cyclophosphamide-induced hematosuppression (Wang et al, 2009).

HSCs differ from the embryonic stem cells in one important aspect – they cannot give rise to the whole spectrum of cell types occurring in the body. The HSC is a member of a family of tissue-specific stem cell, among which are also counted neural stem cells, cord blood stem cell from the umbilical cord, bone marrow stromal stem cells. As the name “tissue-specific stem cells” suggests, these stem cells are only used to replenish specific types of tissues. They are multipotent.

Special places within the bone marrow housing HSCs have been postulated. These are called niches, and currently two distinct types of them are recognised. The niche is a place in which a HSC receives signals that regulate its function or induce dormancy, be it by soluble metabolites, cytokines or direct cell-to-cell contact with non-HSCs forming the niche. Each hematopoietic stem cell can upon division follow either of two routes – (a) maturation through sequential loss of potency and exit from the niche (b) staying in place and continuing with the production of its own clones or entering the state of quiescence.




Figure 6. A simplified overview of hematopoiesis. The self-renewal capacity of HSCs is indicated by the arrow. The gradual loss of colouring indicates increased lineage commitment (Gerrits et al, 2008).

Of the two aforementioned types of niches, the endosteal niche seems to be the place where dormant and slow cycling HSCs with the biggest potential for self-renewal are located. The niche comprises cells of mesenchymal origin which directly interact with the HSCs. The HSCs residing in this niche are called long-term hematopoietic stem cells (LT-HSCs), which reflects their ability contribute to hematopoiesis over a long period of time or even a lifetime.

The microenvironment which is typical for this niche is made up by stromal cells – osteoblasts, adipocytes, endothelial cells and fibroblasts, and the endothelium is in direct contact with the calcified bone matter. A population of specialised N-cadherin-expressing osteoblasts (SNOs) has been found to be absolutely essential for HSCs. These SNOs communicate with the HSCs through homotypic N-cadherin interactions (Zhang et al, 2003). Experiments with mice that had the numbers of bone marrow osteoblasts dramatically lowered (conditional or permanent ablation) showed that these mice had also insufficient numbers of LT-HSCs and, as a direct consequence of that, substantial decrease of lymphoid, erythroid and myeloid

progenitors (including osteoclast progenitors) was observed (Visnjic et al, 2004). The interactions in the niche do not seem to be only one-way, as it has been noted that HSCs drive mesenchymal cells toward osteoblastic lineage when they are grown in a co-culture.

The other distinct type of niche is a vascular one. This niche is supposedly inhabited by short-term hematopoietic stem cells (ST-HSCs) that have migrated here from the endosteal niche. As is to be expected, the architecture and overall microenvironment of the niche differs from the endosteal one. In the vascular niche the HSCs are in contact with the fenestrated endothelium of capillaries, which enables them to closely monitor concentrations of molecules dissolved in blood. Therefore a swift response could be mounted should the situation require it (Yang et al, 2005).

The ST-HSCs give rise to a progeny of multipotent progenitor cells (MPPs). The MPPs can support generation of mature blood cells of all types, but lack the ability to self-renew. The ST-HSCs have in contrast to the LT-HSCs a significantly lower capacity for self-renewal and consequently can single-handedly support hematopoiesis for several weeks only, which is why LT-HSCs are by far better suited for hematopoietic stem cell transplantations. HSCs transplantations are typically carried out in an attempt to correct aberrant hematopoiesis, therefore a long-term solution to the patient's problems is desirable.

Hematopoiesis is complex process, regulation of which is not yet fully clarified, even though a great mass of processes has been studied and our knowledge of it is quite extensive. Apart from direct cell contact and communication between endosteal and vascular niches, cytokines, the endocrine system (regulation of erythropoiesis by erythropoietin) and metabolic factors (such as varied oxygen levels) all take part in it. For example granulocyte colony-stimulating factor (G-CSF), granulocyte-monocyte colony-stimulating factor (GM-CSF), macrophage colony-stimulating factor (M-CSF) are all powerful stimulators driving hematopoietic progenitors to differentiation *in vitro*. However, *in vivo* deficiencies in these factors do not necessarily lead to a certain cell type being completely absent (Lieschke et al, 1994). It seems that these factors do not solely provide lineage-specific signals for

differentiation of HSCs and multipotent progenitors, but rather stimulate the survival/expansion of progenitors, and play a greater role at the terminal maturation stages. On the other hand, interleukin-7 (IL-7) is utterly irreplaceable when it comes to B-cell differentiation. Mice deficient in IL-7 still do produce common lymphoid progenitor cells (CLPs). B-cell production, however, is blocked. The IL-7 regulates the immunoglobulin gene rearrangement once the cell has committed to the B-cell lineage and entered the pro-B-cell stage (von Freeden-Jeffry et al, 1995).

Hematopoietic stem cells are seldom to be seen circulating in the blood (Abkowitz et al, 2003). During the period of embryonic development they migrate from liver and “home” into the bone marrow, establish a stable population in there, and produce high levels of immature and maturing cells of all lineages. The HSCs can be encountered in blood during leukemias and various inflammatory conditions, after injection of granulocyte colony-stimulating factor (G-CSF), and after administration of cyclophosphamide (Szumilas et al, 2005).

3.1. Cyclophosphamide Effects on Hematopoiesis

3.1.1. Myeloablation, Myelosuppression and Immunosuppression

Cyclophosphamide was developed in the 1960s with the objective to create a powerful new antineoplastic agent that would be maximally efficient in battling cancerous growth and, at the same time, be as little toxic to the patients’ healthy tissues as possible. Although it has turned out to be pretty competent in the first field, problems connected to its overall toxicity do occur. As it is the case for virtually any antineoplastic drug and radiation therapy, the regimen affects fast-dividing cells the most – probably due to the DNA being more easily accessible when the nuclear envelope dissolves during mitosis – unfortunately neoplastic tissues are not the only ones which divide fairly quickly. The hematopoietic system housed in the bone marrow is another great target for anti-cancer therapeutics, with hematopoietic progenitor cells being the most vulnerable (DeWys et al, 1970; Valeriote et al, 1968). Severe Myelosuppression is to this day one of the key dose-limiting aspects of cyclophosphamide administration in the clinical setting, and might even happen to be a reason for discontinuing chemotherapy in some patients (Presant & Klahr, 1977).

Injury to the bone marrow by a cyclophosphamide-containing regimen of treatment usually leads to a pronounced decline in the granulocyte, lymphocyte, thrombocyte numbers (Fried & Johnson, 1968; Mosienko et al, 2002), and severe cases of anemia, which is in accordance with an observable decrease in bone marrow cellularity (Sefc et al, 2003). While the cases of developed anemia and thrombocytopenia can be confronted by administering blood and thrombocyte transfusions respectively, the transfusions are still just a way of minimising the damage without solving the underlying problem.

The most serious of all cytopenias induced by cyclophosphamide is neutropenia. Patients who see their neutrophil levels fall below the value of 100 cells/mm³ (severe neutropenia) are extremely prone to infection by pathogens such as *Klebsiella pneumoniae*, and are incapable of initiating an efficient immune response against them. A concomitant administration of prophylactic antibiotic drugs is therefore required (Tjan-Heijnen et al, 2001). Treatment protocols also sometimes include G-CSF as it has been shown that it can reduce neutropenia (Neidhart et al, 1989).

In a stark contrast to the undesired chemotherapy-induced immunosuppression in cancer sufferers is the intentional prescription of cyclophosphamide which aims to achieve this result. Cyclophosphamide is a drug of choice for the management of some very severe immune disorders (Burt et al, 1998). Among these are systemic lupus erythematosus (Katsifis et al, 2002), systemic sclerosis; and cyclophosphamide is currently being tested as a promising therapy for autoimmune aplastic anemia patients (Brodsky et al, 2004; Brodsky et al, 2001).

Yet another situation in which immunosuppression by cyclophosphamide is desirable is the pre-transplantation conditioning of the intended host (Kapoor et al, 1981; Litzow et al, 2002). Elimination of host alloreactive lymphocytes serves as a prevention of the rejection of a transplant (Aversa et al, 1999). Moreover, high-dose cyclophosphamide seems to be quite efficient even in the capacity of single-agent prophylaxis against both acute and chronic graft-versus-host disease (Luznik et al).

The myelosuppressive effect has been attributed to cyclophosphamide right from the very beginning of its use in cancer therapy. Mice that were subjected to cyclophosphamide in doses equivalent of bone marrow-injury causing γ -irradiation in early studies had been found to develop a corresponding response of a trauma and subsequent improvement. What was more of an interest was the fact that murine bone marrow and peripheral blood cell count recovered more swiftly after cyclophosphamide regimen. Moreover an overshoot in reticulocytes could be observed – so-called rebound reticulocytosis (Valeriote et al, 1968). While the myelotoxic effect of cyclophosphamide has an unquestionably great negative impact on experimental models (mostly mice) and human patients both, it is nonetheless only temporary and can be weathered with supportive therapy.

The transient nature of cyclophosphamide-induced myelosuppression is attributable to its mode of action. While rapidly proliferating hematopoietic progenitor cells are hit with full force, slow-cycling hematopoietic stem cells are more resistant to the drug. The conservation of HSCs is intrinsic for hematopoietic reconstitution. Cyclophosphamide regimens need to be established so that HSCs have enough time to replenish the pool of HPCs. During the period of hematopoietic reconstitution, when the HSCs divide rapidly to compensate the loss of progenitor and mature blood cells, are even these cells susceptible to damage by cyclophosphamide, which is counterproductive in regimens where the HSCs are not the intended targets.

In recent years, various studies have been focused on eliminating, or at least decreasing, the hematologic toxicity of chemotherapeutic agents. Notably macrophage inflammatory protein-1 alpha (MIP-1 α) piqued researchers' interest due to its ability to cause cell-cycle arrest, which could serve as protection for HPCs. However, this chemokine naturally polymerises in extremely low concentrations (>20 ng/ml), which consequently renders it inactive. Retaining its native form by means of organic solvents was not a suitable option for patient therapy. A recombinant protein with significantly lowered polymerising tendencies has been developed. While it cannot prevent cyclophosphamide-induced neutropenia or its

severity, it seems to speed up the recovery process, which is promising for the future (Gilmore et al, 1999; Marshall et al, 1997).

Meanwhile cyclophosphamide serves as an effective means of making murine models immunosuppressed, which is very useful for exact evaluation of the efficacy of antibiotics (Zuluaga et al, 2006).

3.1.2. Mobilisation of Hematopoietic Stem Cells

The list of effects cyclophosphamide has on hematopoiesis by no means ends at myelosuppression and immunosuppression. Another important outcome of cyclophosphamide administration is the mobilisation of hematopoietic stem and progenitor cells (sorted according to the expression of surface CD34⁺ marker) into the peripheral blood. This is widely used in clinical practice for mobilisation of CD34⁺ cells into the peripheral blood, from where they can be collected by a process of apheresis and further used for autologous transplantations (Rosen et al, 2000). Mobilisation protocols do not usually employ cyclophosphamide as the sole agent as it has been demonstrated that significantly higher numbers of progenitors are collected when G-CSF or GM-CSF are included in the regimen (Caporali et al, 2001). The aforementioned growth factors can also stimulate mobilisation singlehandedly (Abkowitz et al, 2003), but their combination with chemotherapy is also beneficial in terms of yield. Nevertheless, cyclophosphamide is not included in mobilisation regimens in healthy donors due to its toxic properties, because it would simply not be ethical to impose those on healthy individuals.

The process of release of HSCs and HPCs from the bone marrow into the peripheral blood has not yet been completely elucidated, although some important discoveries have been made. Being de facto the direct opposite of homing, the attention of researches has been turned to adhesive molecular pairs such as very late antigen 4 (VLA-4) and vascular cell adhesion molecule-1 [VCAM-1 (Kikuta et al, 2000)]; and chemokines and their receptors such as stromal cell-derived factor 1 (SDF-1/CXCL12) and its receptor CXCR4. The evidence for the importance of VLA-4/VCAM-1 stems from studies utilising anti-VLA-4 antibodies. The administration of VLA-4 antibodies leads to successful mobilisation of HSCs in

experimental animals (Papayannopoulou & Nakamoto, 1993). Neutralisation of VLA-4 also prevents homing of transplanted HSCs (Zanjani et al, 1999).

SDF-1 is highly expressed in the bone marrow, notably by the immature osteoblasts of the endosteal niche. The gradient of SDF-1 is deemed necessary for homing of CD34⁺ both in steady-state hematopoiesis and after transplantation into myeloablated hosts. CXCR4 is a membrane-bound receptor protein with seven transmembrane domains. It is linked to G-proteins and the signal cascade initiated upon interaction with SDF-1 may result for example in the reorganisation of actin cytoskeleton and formation of pseudopodia. SDF-1 α induces phosphorylation of multiple adhesion proteins. The importance of SDF-1/CXCR4 axis for hematopoiesis has been demonstrated *in vivo* when knock-out mice lacking both molecules developed severe defects in B lymphopoiesis and myelopoiesis.

On the other hand, mobilisation can be induced in mice by artificial overexpression of SDF-1, as too high a concentration leads to desensitisation and internalisation/degradation of CXCR4 of the target cells and thus to their unresponsiveness to SDF-1 (Alsayed et al, 2007). This discovery was most important as it had been previously shown that chemotherapy tends to increase the levels of SDF-1 in the bone marrow, and yet cyclophosphamide does facilitate mobilisation, which naturally seemed contradictory. Moreover, so-called “poor mobilisers” in whom achievement of target numbers of collected CD34⁺ cells for autologous transplantations is problematic do express more SDF-1 at steady state than “good mobilisers” and healthy individuals (Gazitt & Liu, 2001). Even though mobilised HSCs express significantly less CXCR4 (Carion et al, 2003; Gazitt & Liu, 2001), they are able to home into the hosts bone marrow. This is attributed to chemokine and overall environment of the hosts circulatory system which promotes the production of new CXCR4 receptors on the cell surface.

It has been observed that rising numbers of hematopoietic progenitors in the peripheral blood are in direct correlation with increased levels of proteolytic enzymes in the bone marrow (Levesque et al, 2002). Matrix metalloproteinase 9 (Carion et al, 2003), cathepsin G and leucocyte elastase all degrade the N-terminus of

SDF-1 *in vitro* which disallows binding to the receptor and results in the loss of its chemoattractant function.

AMD-3100 (also known as plerixafor) is a substance causing reversible inhibition of CXCR4 (Hendrix et al, 2000). When it is combined with G-CSF in human donors for mobilisation, the collected cells express higher levels of VLA-4, tend to migrate to SDF-1 better, and also show significantly better engraftment in NOD/SCID (non-obese diabetic/severe combined immunodeficiency) hosts. This example of crosstalk between chemokines and/or adhesion molecules is not the only of the kind, and much effort is currently being put into deciphering the relationship status of various chemokine/adhesion molecule interaction-induced responses of HSCs (Peled et al, 2000).

4. Conclusion

Cyclophosphamide is an old drug put to new uses as we gradually discover how a human body works. Its precise metabolism and mode of action has long been a mystery, but decades of research are slowly heading to illumination of the principles.

It is now possible to fully appreciate how a polymorphism in a gene coding a particular enzyme influences the metabolism of the drug, and interpatient differences can be addressed. As the mankind sees the numbers of malignancies rising, it is most beneficial that cyclophosphamide's mode of action is better understood than it used to be fifty years ago.

Its use in hematology has been extensively studied. Interestingly, cyclophosphamide can mimic both the impact of γ -irradiation and of the hematopoietic growth factor G-CSF depending on dosage, timing and genetic background of the subject.

In its capacity of mobilisation-inducing drug cyclophosphamide demonstrated quite similar tendencies to G-CSF – after its administration, bone marrow environment reacts by releasing stored neutrophil proteases which cleave molecules facilitating chemotaxis and adhesion to the bone marrow niche, thus enabling the egress of stem and progenitor cells into the peripheral blood. Among the affected molecular pairs studied in recent years are CXCR4 and its ligand SDF-1, and VLA-4 with VCAM-1. These interactions cannot be viewed as isolated systems, as one stimulus can often influence the way another one is processed, and there is significant amount of cross-talk going on. More studies of these kinds of interactions are yet to come and will hopefully lead to improvement of clinical protocols and brighter outlook for patients with hematologic and immune malignancies, whose numbers are rapidly growing.

5. References:

- Abkowitz JL, Robinson AE, Kale S, Long MW, Chen J (2003) Mobilization of hematopoietic stem cells during homeostasis and after cytokine exposure. *Blood* **102**(4): 1249-1253
- Alsayed Y, Ngo H, Runnels J, Leleu X, Singha UK, Pitsillides CM, Spencer JA, Kimlinger T, Ghobrial JM, Jia X, Lu G, Timm M, Kumar A, Cote D, Veilleux I, Hedin KE, Roodman GD, Witzig TE, Kung AL, Hideshima T, Anderson KC, Lin CP, Ghobrial IM (2007) Mechanisms of regulation of CXCR4/SDF-1 (CXCL12)-dependent migration and homing in multiple myeloma. *Blood* **109**(7): 2708-2717
- Aversa F, Terenzi A, Carotti A, Felicini R, Jacucci R, Zei T, Latini P, Aristei C, Santucci A, Martelli MP, Cunningham I, Reisner Y, Martelli MF (1999) Improved outcome with T-cell-depleted bone marrow transplantation for acute leukemia. *J Clin Oncol* **17**(5): 1545-1550
- Brodsky RA, Chen AR, Brodsky I, Jones RJ (2004) High-dose cyclophosphamide as salvage therapy for severe aplastic anemia. *Exp Hematol* **32**(5): 435-440
- Brodsky RA, Sensenbrenner LL, Smith BD, Dorr D, Seaman PJ, Lee SM, Karp JE, Brodsky I, Jones RJ (2001) Durable treatment-free remission after high-dose cyclophosphamide therapy for previously untreated severe aplastic anemia. *Ann Intern Med* **135**(7): 477-483
- Burt RK, Traynor AE, Pope R, Schroeder J, Cohen B, Karlin KH, Lobeck L, Goolsby C, Rowlings P, Davis FA, Stefoski D, Terry C, Keever-Taylor C, Rosen S, Vesole D, Fishman M, Brush M, Mujias S, Villa M, Burns WH (1998) Treatment of autoimmune disease by intense immunosuppressive conditioning and autologous hematopoietic stem cell transplantation. *Blood* **92**(10): 3505-3514
- Caporali R, Perotti C, Pedrazzoli P, Da Prada GA, Bernuzzi S, Montecucco C (2001) Cyclophosphamide plus granulocyte colony stimulating factor (G-CSF) is more effective than G-CSF alone in mobilizing hemopoietic progenitor cells in severe, refractory rheumatoid arthritis. *Haematologica* **86**(1): 106-107
- Carion A, Benboubker L, Herault O, Roingeard F, Degenne M, Senecal D, Desbois I, Colombat P, Charbord P, Binet C, Domenech J (2003) Stromal-derived factor 1 and matrix metalloproteinase 9 levels in bone marrow and peripheral blood of patients mobilized by granulocyte colony-stimulating factor and chemotherapy. Relationship with mobilizing capacity of haematopoietic progenitor cells. *Br J Haematol* **122**(6): 918-926
- Crook TR, Souhami RL, McLean AE (1986) Cytotoxicity, DNA cross-linking, and single strand breaks induced by activated cyclophosphamide and acrolein in human leukemia cells. *Cancer Res* **46**(10): 5029-5034

de Jonge ME, Huitema AD, Rodenhuis S, Beijnen JH (2005) Clinical pharmacokinetics of cyclophosphamide. *Clin Pharmacokinet* **44**(11): 1135-1164

de Montellano PRO (2005) *Cytochrome P450: Structure, Mechanism, and Biochemistry*, 3rd edn.: Springer US.

DeWys WD, Goldin A, Man, El N (1970) Hematopoietic recovery after large doses of cyclophosphamide: correlation of proliferative state with sensitivity. *Cancer Res* **30**(6): 1692-1697

Ellis J, Gutierrez A, Barsukov IL, Huang WC, Grossmann JG, Roberts GC (2009) Domain motion in cytochrome P450 reductase: conformational equilibria revealed by NMR and small-angle x-ray scattering. *J Biol Chem* **284**(52): 36628-36637

Fried W, Johnson C (1968) The effect of cyclophosphamide on hematopoietic stem cells. *Radiat Res* **36**(3): 521-527

Gazitt Y, Liu Q (2001) Plasma levels of SDF-1 and expression of SDF-1 receptor on CD34+ cells in mobilized peripheral blood of non-Hodgkin's lymphoma patients. *Stem Cells* **19**(1): 37-45

Gerrits A, Dykstra B, Otten M, Bystrykh L, de Haan G (2008) Combining transcriptional profiling and genetic linkage analysis to uncover gene networks operating in hematopoietic stem cells and their progeny. *Immunogenetics* **60**(8): 411-422

Gilmore GL, DePasquale DK, Shadduck RK (1999) Protective effects of BB-10010 treatment on chemotherapy-induced neutropenia in mice. *Exp Hematol* **27**(2): 195-202

Griskevicius L, Yasar U, Sandberg M, Hidestrand M, Eliasson E, Tybring G, Hassan M, Dahl ML (2003) Bioactivation of cyclophosphamide: the role of polymorphic CYP2C enzymes. *Eur J Clin Pharmacol* **59**(2): 103-109

Gu J, Chen CS, Wei Y, Fang C, Xie F, Kannan K, Yang W, Waxman DJ, Ding X (2007) A mouse model with liver-specific deletion and global suppression of the NADPH-cytochrome P450 reductase gene: characterization and utility for in vivo studies of cyclophosphamide disposition. *J Pharmacol Exp Ther* **321**(1): 9-17

Gurtoo HL, Hipkens JH, Sharma SD (1981) Role of glutathione in the metabolism-dependent toxicity and chemotherapy of cyclophosphamide. *Cancer Res* **41**(9 Pt 1): 3584-3591

Harleton E, Webster M, Bumpus NN, Kent UM, Rae JM, Hollenberg PF (2004) Metabolism of N,N',N''-triethylenethiophosphoramidate by CYP2B1 and CYP2B6

results in the inactivation of both isoforms by two distinct mechanisms. *J Pharmacol Exp Ther* **310**(3): 1011-1019

Hendrix CW, Flexner C, MacFarland RT, Giandomenico C, Fuchs EJ, Redpath E, Bridger G, Henson GW (2000) Pharmacokinetics and safety of AMD-3100, a novel antagonist of the CXCR-4 chemokine receptor, in human volunteers. *Antimicrob Agents Chemother* **44**(6): 1667-1673

Henry CJ, Brewer WG, Jr., Royer NS (1994) Hematological and clinical responses to combined mitoxantrone and cyclophosphamide administration to normal cats. *Can Vet J* **35**(11): 706-708

Highley MS, Schrijvers D, Van Oosterom AT, Harper PG, Momerency G, Van Cauwenberghe K, Maes RA, De Bruijn EA, Edelstein MB (1997) Activated oxazaphosphorines are transported predominantly by erythrocytes. *Ann Oncol* **8**(11): 1139-1144

Huang Z, Roy P, Waxman DJ (2000) Role of human liver microsomal CYP3A4 and CYP2B6 in catalyzing N-dechloroethylation of cyclophosphamide and ifosfamide. *Biochem Pharmacol* **59**(8): 961-972

Chang TK, Yu L, Maurel P, Waxman DJ (1997) Enhanced cyclophosphamide and ifosfamide activation in primary human hepatocyte cultures: response to cytochrome P-450 inducers and autoinduction by oxazaphosphorines. *Cancer Res* **57**(10): 1946-1954

Kapoor N, Kirkpatrick D, Blaese RM, Oleske J, Hilgartner MH, Chaganti RS, Good RA, O'Reilly RJ (1981) Reconstitution of normal megakaryocytopoiesis and immunologic functions in Wiskott-Aldrich syndrome by marrow transplantation following myeloablation and immunosuppression with busulfan and cyclophosphamide. *Blood* **57**(4): 692-696

Katsifis GE, Tzioufas AG, Vlachoyiannopoulos PG, Voulgarelis M, Moutsopoulos HM, Ioannidis JP (2002) Risk of myelotoxicity with intravenous cyclophosphamide in patients with systemic lupus erythematosus. *Rheumatology (Oxford)* **41**(7): 780-786

Kenney LB, Laufer MR, Grant FD, Grier H, Diller L (2001) High risk of infertility and long term gonadal damage in males treated with high dose cyclophosphamide for sarcoma during childhood. *Cancer* **91**(3): 613-621

Kikuta T, Shimazaki C, Ashihara E, Sudo Y, Hirai H, Sumikuma T, Yamagata N, Inaba T, Fujita N, Kina T, Nakagawa M (2000) Mobilization of hematopoietic primitive and committed progenitor cells into blood in mice by anti-vascular adhesion molecule-1 antibody alone or in combination with granulocyte colony-stimulating factor. *Exp Hematol* **28**(3): 311-317

- Kohn KW, Hartley JA, Mattes WB (1987) Mechanisms of DNA sequence selective alkylation of guanine-N7 positions by nitrogen mustards. *Nucleic Acids Res* **15**(24): 10531-10549
- Kwon CH, Maddison K, LoCastro L, Borch RF (1987) Accelerated decomposition of 4-hydroxycyclophosphamide by human serum albumin. *Cancer Res* **47**(6): 1505-1508
- Lawrence HJ, Simone J, Aur RJ (1975) Cyclophosphamide-induced hemorrhagic cystitis in children with leukemia. *Cancer* **36**(5): 1572-1576
- LeBlanc GA, Waxman DJ (1990) Mechanisms of cyclophosphamide action on hepatic P-450 expression. *Cancer Res* **50**(18): 5720-5726
- Levesque JP, Hendy J, Takamatsu Y, Williams B, Winkler IG, Simmons PJ (2002) Mobilization by either cyclophosphamide or granulocyte colony-stimulating factor transforms the bone marrow into a highly proteolytic environment. *Exp Hematol* **30**(5): 440-449
- Lieschke GJ, Stanley E, Grail D, Hodgson G, Sinickas V, Gall JA, Sinclair RA, Dunn AR (1994) Mice lacking both macrophage- and granulocyte-macrophage colony-stimulating factor have macrophages and coexistent osteopetrosis and severe lung disease. *Blood* **84**(1): 27-35
- Litzow MR, Perez WS, Klein JP, Bolwell BJ, Camitta B, Copelan EA, Gale RP, Giralt SA, Keating A, Lazarus HM, Marks DI, McCarthy PL, Miller CB, Milone G, Prentice HG, Russell JA, Schultz KR, Trigg ME, Weisdorf DJ, Horowitz MM (2002) Comparison of outcome following allogeneic bone marrow transplantation with cyclophosphamide-total body irradiation versus busulphan-cyclophosphamide conditioning regimens for acute myelogenous leukaemia in first remission. *Br J Haematol* **119**(4): 1115-1124
- Luznik L, Bolanos-Meade J, Zahurak M, Chen AR, Smith BD, Brodsky R, Huff CA, Borrello I, Matsui W, Powell JD, Kasamon Y, Goodman SN, Hess A, Levitsky HI, Ambinder RF, Jones RJ, Fuchs EJ High-dose cyclophosphamide as single-agent, short-course prophylaxis of graft-versus-host disease. *Blood* **115**(16): 3224-3230
- Manthey CL, Landkamer GJ, Sladek NE (1990) Identification of the mouse aldehyde dehydrogenases important in aldophosphamide detoxification. *Cancer Res* **50**(16): 4991-5002
- Marshall E, Woolford LB, Lord BI (1997) Continuous infusion of macrophage inflammatory protein MIP-1 α enhances leucocyte recovery and haemopoietic progenitor cell mobilization after cyclophosphamide. *Br J Cancer* **75**(12): 1715-1720
- Meunier B, de Visser SP, Shaik S (2004) Mechanism of oxidation reactions catalyzed by cytochrome p450 enzymes. *Chem Rev* **104**(9): 3947-3980

Mosienko MD, Lyniv LS, Kireeva SS, Ryabukha VM, Mosienko VS (2002) The Protective Action of Immunomodulator of Bacterial Origin and Melatonin in Mice with Cyclophosphamide-induced Myelosuppression. *Experimental Oncology* **24**(2): 145-149

Neidhart J, Mangalik A, Kohler W, Stidley C, Saiki J, Duncan P, Souza L, Downing M (1989) Granulocyte colony-stimulating factor stimulates recovery of granulocytes in patients receiving dose-intensive chemotherapy without bone marrow transplantation. *J Clin Oncol* **7**(11): 1685-1692

Papayannopoulou T, Nakamoto B (1993) Peripheralization of hemopoietic progenitors in primates treated with anti-VLA4 integrin. *Proc Natl Acad Sci U S A* **90**(20): 9374-9378

Pass GJ, Carrie D, Boylan M, Lorimore S, Wright E, Houston B, Henderson CJ, Wolf CR (2005) Role of hepatic cytochrome p450s in the pharmacokinetics and toxicity of cyclophosphamide: studies with the hepatic cytochrome p450 reductase null mouse. *Cancer Res* **65**(10): 4211-4217

Peled A, Kollet O, Ponomarev T, Petit I, Franitza S, Grabovsky V, Slav MM, Nagler A, Lider O, Alon R, Zipori D, Lapidot T (2000) The chemokine SDF-1 activates the integrins LFA-1, VLA-4, and VLA-5 on immature human CD34(+) cells: role in transendothelial/stromal migration and engraftment of NOD/SCID mice. *Blood* **95**(11): 3289-3296

Phillips AH, Langdon RG (1962) Hepatic triphosphopyridine nucleotide-cytochrome c reductase: isolation, characterization, and kinetic studies. *J Biol Chem* **237**: 2652-2660

Presant CA, Klahr C (1977) Severe myelosuppression from piperazinedione, (NSC No. 135758), cyclophosphamide plus dimethyl-triazeno-imidazole carboxamide (DTIC). *Cancer* **40**(4): 1386-1389

Rae JM, Soukhova NV, Flockhart DA, Desta Z (2002) Triethylenethiophosphoramide is a specific inhibitor of cytochrome P450 2B6: implications for cyclophosphamide metabolism. *Drug Metab Dispos* **30**(5): 525-530

Ren S, Yang JS, Kalhorn TF, Slattery JT (1997) Oxidation of cyclophosphamide to 4-hydroxycyclophosphamide and deschloroethylcyclophosphamide in human liver microsomes. *Cancer Res* **57**(19): 4229-4235

Rosen O, Thiel A, Massenkeil G, Hiepe F, Haupl T, Radtke H, Burmester GR, Gromnica-Ihle E, Radbruch A, Arnold R (2000) Autologous stem-cell transplantation in refractory autoimmune diseases after in vivo immunoablation and ex vivo depletion of mononuclear cells. *Arthritis Res* **2**(4): 327-336

- Roy P, Yu LJ, Crespi CL, Waxman DJ (1999) Development of a substrate-activity based approach to identify the major human liver P-450 catalysts of cyclophosphamide and ifosfamide activation based on cDNA-expressed activities and liver microsomal P-450 profiles. *Drug Metab Dispos* **27**(6): 655-666
- Sefc L, Psenak O, Sykora V, Sulc K, Necas E (2003) Response of hematopoiesis to cyclophosphamide follows highly specific patterns in bone marrow and spleen. *J Hematother Stem Cell Res* **12**(1): 47-61
- Singer A, Maldjian P, Simmons MZ (2004) Extramedullary hematopoiesis presenting as a focal splenic mass: a case report. *Abdom Imaging* **29**(6): 710-712
- Struck RF, Alberts DS, Horne K, Phillips JG, Peng YM, Roe DJ (1987) Plasma pharmacokinetics of cyclophosphamide and its cytotoxic metabolites after intravenous versus oral administration in a randomized, crossover trial. *Cancer Res* **47**(10): 2723-2726
- Szumilas P, Barcew K, Baskiewicz-Masiuk M, Wiszniewska B, Ratajczak MZ, Machalinski B (2005) Effect of stem cell mobilization with cyclophosphamide plus granulocyte colony-stimulating factor on morphology of haematopoietic organs in mice. *Cell Prolif* **38**(1): 47-61
- Tjan-Heijnen VC, Postmus PE, Ardizzoni A, Manegold CH, Burghouts J, van Meerbeeck J, Gans S, Mollers M, Buchholz E, Biesma B, Legrand C, Debruyne C, Giaccone G (2001) Reduction of chemotherapy-induced febrile leucopenia by prophylactic use of ciprofloxacin and roxithromycin in small-cell lung cancer patients: an EORTC double-blind placebo-controlled phase III study. *Ann Oncol* **12**(10): 1359-1368
- Valeriote FA, Collins DC, Bruce WR (1968) Hematological recovery in the mouse following single doses of gamma radiation and cyclophosphamide. *Radiat Res* **33**(3): 501-511
- Vermilion JL, Ballou DP, Massey V, Coon MJ (1981) Separate roles for FMN and FAD in catalysis by liver microsomal NADPH-cytochrome P-450 reductase. *J Biol Chem* **256**(1): 266-277
- Visnjic D, Kalajzic Z, Rowe DW, Katavic V, Lorenzo J, Aguila HL (2004) Hematopoiesis is severely altered in mice with an induced osteoblast deficiency. *Blood* **103**(9): 3258-3264
- von Freeden-Jeffry U, Vieira P, Lucian LA, McNeil T, Burdach SE, Murray R (1995) Lymphopenia in interleukin (IL)-7 gene-deleted mice identifies IL-7 as a nonredundant cytokine. *J Exp Med* **181**(4): 1519-1526

Wang Y, Meng Q, Qiao H, Jiang H, Sun X (2009) Role of the spleen in cyclophosphamide-induced hematosuppression and extramedullary hematopoiesis in mice. *Arch Med Res* **40**(4): 249-255

Xie HJ, Yasar U, Lundgren S, Griskevicius L, Terelius Y, Hassan M, Rane A (2003) Role of polymorphic human CYP2B6 in cyclophosphamide bioactivation. *Pharmacogenomics J* **3**(1): 53-61

Yang L, Bryder D, Adolfsson J, Nygren J, Mansson R, Sigvardsson M, Jacobsen SE (2005) Identification of Lin(-)Sca1(+)kit(+)CD34(+)Flt3- short-term hematopoietic stem cells capable of rapidly reconstituting and rescuing myeloablated transplant recipients. *Blood* **105**(7): 2717-2723

Zanjani ED, Flake AW, Almeida-Porada G, Tran N, Papayannopoulou T (1999) Homing of human cells in the fetal sheep model: modulation by antibodies activating or inhibiting very late activation antigen-4-dependent function. *Blood* **94**(7): 2515-2522

Zhang J, Niu C, Ye L, Huang H, He X, Tong WG, Ross J, Haug J, Johnson T, Feng JQ, Harris S, Wiedemann LM, Mishina Y, Li L (2003) Identification of the haematopoietic stem cell niche and control of the niche size. *Nature* **425**(6960): 836-841

Zuluaga AF, Salazar BE, Rodriguez CA, Zapata AX, Agudelo M, Vesga O (2006) Neutropenia induced in outbred mice by a simplified low-dose cyclophosphamide regimen: characterization and applicability to diverse experimental models of infectious diseases. *BMC Infect Dis* **6**: 55